GENERAL

nanda nursing diagnosis for chronic kidney disease

Published: September 3, 2026 | Index ID: #-

Introduction

Chronic kidney disease (CKD) is a progressive condition that gradually diminishes kidney function, leading to a host of metabolic, cardiovascular, and psychosocial complications. As the prevalence of CKD rises worldwide, nurses play a pivotal role in early detection, comprehensive assessment, and targeted care planning. Central to this process is the use of the North American Nursing Diagnosis Association (NANDA) taxonomy, which provides a standardized language for identifying patient problems and guiding interventions.

In this article, we will explore **nanda nursing diagnosis for chronic kidney disease**in depth. We'll examine the core concepts of CKD, the NANDA framework, and how to apply specific nursing diagnoses to individual patients. Practical assessment tools, evidence based interventions, and outcome measures will be highlighted to equip nurses, educators, and healthcare teams with the knowledge needed to deliver high quality, patient centered care.

Understanding Chronic Kidney Disease

What Is CKD?

CKD refers to the gradual loss of kidney function over months or years. It is classified into five stages based on the estimated glomerular filtration rate (eGFR) and albuminuria levels:

- Stage 1: eGFR "e90 /mL/min/1.73 /m² with kidney damage
- Stage 2: eGFR 60–89 /mL/min/1.73 /m²
- Stage 3a: eGFR 45–59 /mL/min/1.73 /m²
- Stage 3b: eGFR 30–44 /mL/min/1.73 /m²
- Stage 4: eGFR 15–29 /mL/min/1.73 /m²
- Stage 5: eGFR

Common causes include diabetes mellitus, hypertension, glomerulonephritis, polycystic kidney disease, and prolonged use of nephrotoxic medications. CKD is often silent in early stages, which underscores the importance of routine screening in high risk populations.

Clinical Manifestations

As CKD progresses, patients may experience:

- Edema (especially lower extremities)
- Hypertension or uncontrolled blood pressure
- Fatigue, weakness, and decreased appetite
- Electrolyte disturbances (hyperkalemia, hyperphosphatemia)
- Acid–base imbalance (metabolic acidosis)
- Anemia and bone mineral disorders
- Psychosocial stress, depression, or anxiety

These manifestations provide the clinical clues that nurses use to formulate accurate NANDA diagnoses.

The NANDA Framework

Why Use NANDA?

The North American Nursing Diagnosis Association taxonomy offers a structured, evidence based language that:

- Facilitates clear communication among interdisciplinary teams
- Supports outcome measurement and quality improvement
- Enhances documentation for legal and billing purposes
- Promotes consistent, patient centered care planning

By applying NANDA diagnoses to CKD, nurses can identify both the physiological and psychosocial aspects of the disease, ensuring comprehensive care.

Key Components of a NANDA Diagnosis

1. Problem Statement: The patient's actual or potential health issue.
2. Defining Characteristics: Observable signs or measurable data that confirm the diagnosis.
3. Related Factors: Underlying causes or contributing factors.
4. Risk Factors: Conditions that increase the likelihood of developing the problem.

Common NANDA Nursing Diagnoses for CKD

Below is a curated list of NANDA diagnoses frequently encountered in patients with chronic kidney disease. Each diagnosis is accompanied by defining characteristics, related factors, and suggested interventions.

1. Fluid Volume Excess

- Defining Characteristics: Edema, weight gain, elevated blood pressure, jugular venous distension.
- Related Factors: Reduced renal excretion, sodium retention, impaired diuretic response.
- Interventions: Monitor intake/output, administer diuretics as ordered, restrict sodium, encourage fluid balance education.

2. Fluid Volume Deficit

- Defining Characteristics: Dry mucous membranes, decreased skin turgor, orthostatic hypotension.
- Related Factors: Diuretic therapy, vomiting, diarrhea, inadequate fluid intake.
- Interventions: Assess hydration status, administer IV fluids per protocol, educate on oral rehydration strategies.

3. Risk for Electrolyte Imbalance

- Defining Characteristics: Abnormal serum potassium, sodium, calcium, or phosphate levels.
- Related Factors: Declining GFR, dietary restrictions, medication effects.
- Interventions: Monitor serum electrolytes, administer supplements or binders, coordinate with dietitian for dietary planning.

4. Risk for Infection

- Defining Characteristics: Immunosuppression, catheter use, dialysis access sites.
- Related Factors: Reduced neutrophil function, frequent vascular access, comorbid conditions.
- Interventions: Strict aseptic technique, monitor for signs of infection, administer prophylactic antibiotics when indicated.

5. Impaired Renal Function

- Defining Characteristics: Elevated creatinine, reduced eGFR, proteinuria.
- Related Factors: Chronic hypertension, diabetes, nephrotoxic drugs.
- Interventions: Adjust medication dosing, monitor renal labs, collaborate with nephrologist.

6. Risk for Altered Nutrition: Less Than Body Requirements

- Defining Characteristics: Weight loss, decreased appetite, malnutrition indices.
- Related Factors: Anorexia, metabolic acidosis, dietary restrictions.
- Interventions: Nutritional assessment, dietitian referral, monitor weight and laboratory markers.

7. Ineffective Coping

- Defining Characteristics: Anxiety, depression, refusal of treatment.
- Related Factors: Chronic illness burden, dialysis schedules, fear of complications.
- Interventions: Provide emotional support, involve counseling services, encourage family participation.

Assessment Strategies for CKD Nursing Diagnoses

Accurate assessment is the cornerstone of effective nursing diagnosis. The following tools and techniques help nurses gather comprehensive data:

Vital Sign Monitoring

Track blood pressure, heart rate, respiratory rate, and temperature regularly. Hypertension is a key risk factor for CKD progression.

Laboratory Data Review

Key labs include serum creatinine, eGFR, BUN, electrolytes, hemoglobin, albumin, and urine albumin-to-creatinine ratio. Trends over time provide insight into disease trajectory.

Physical Examination

Assess for edema (pitting), skin turgor, mucous membrane moisture, and jugular venous pressure. Evaluate for signs of fluid overload or deficit.

Patient History and Psychosocial Assessment

Document comorbid conditions, medication history (especially nephrotoxic agents), dietary habits, and psychosocial stressors. Understanding the patient"s perspective aids in selecting appropriate interventions.

Functional Status Evaluation

Use scales such as the Barthel Index or Karnofsky Performance Status to gauge independence and inform care plans.

Intervention Planning: Translating Diagnosis into Action

Once the NANDA diagnoses are identified, the next step is to formulate a comprehensive care plan. Below is a template illustrating how to structure interventions, desired outcomes, and evaluation criteria.

Example Care Plan: Fluid Volume Excess

- Goal: Maintain euvolemia and prevent edema.
- Interventions:

  Assess intake and output every shift.
- Administer diuretics per physician order.
- Restrict sodium to
- Educate patient on low sodium diet and fluid restrictions.
- Monitor orthostatic vital signs.

1. Stable weight within 0.5 /kg of baseline.
2. No pitting edema in lower extremities.
3. Blood pressure

Documentation and Legal Considerations

Accurate documentation of NANDA diagnoses and care plans is essential for continuity of care, legal protection, and reimbursement. Key elements include:

- Clear identification of the diagnosis using the NANDA code.
- Detailed assessment data supporting the diagnosis.
- Specific, measurable outcomes and intervention timelines.
- Regular progress notes documenting changes in status.
- Interdisciplinary collaboration notes (e.g., nephrology consults, dietitian plans).

Case Study: Applying NANDA Diagnoses in a CKD Patient

Mrs. A, a 68 year old woman with type 2 diabetes, presents with Stage 4 CKD (eGFR 22 mL/min/1.73 m^2). She reports fatigue, swelling in her ankles, and decreased appetite. Her laboratory results show hyperkalemia (K z /5.8 /mmol/L) and anemia (Hb /9.5 /g/dL).

Using the NANDA framework, the nursing team identifies the following diagnoses:

- Fluid Volume Excess – related to decreased renal excretion.
- Risk for Electrolyte Imbalance – related to declining GFR.
- Impaired Renal Function – related to diabetic nephropathy.
- Risk for Altered Nutrition – less than body requirements.

Interventions include diuretic therapy, potassium binding resin, dietary counseling, and referral to a dietitian. Outcomes are tracked over four weeks, showing reduced ankle edema, normalized potassium levels, and improved hemoglobin. The case exemplifies how systematic application of NANDA diagnoses leads to measurable improvements.

Evidence Based Interventions for CKD Nursing Diagnoses

Recent research underscores several best practices that nurses can incorporate into their CKD care plans:

1. Sodium Restriction

Studies demonstrate that limiting sodium to

2. Phosphate Binders

Early initiation of phosphate binders helps maintain serum phosphate within normal limits, decreasing cardiovascular risk.

3. Anemia Management

Targeting hemoglobin levels between 10–12 g/dL with erythropoiesis stimulating agents improves quality of life and reduces hospitalization.

4. Early Referral for Dialysis Planning

Initiating discussions about modality options (hemodialysis, peritoneal dialysis, transplant) when eGFR falls below 15 /mL/min/1.73 /m^2 improves patient preparedness and outcomes.

5. Psychosocial Support

Integrating counseling and support groups reduces depression and enhances adherence to treatment plans.

Integrating Technology into CKD Care

Digital tools can streamline assessment and monitoring:

Baca Juga (Artikel Terkait)

- **advanced engineering electromagnetics solution manual edition hardcover by balanis constantine a published by wiley**
  URL: http://denniswildberger.com/advanced-engineering-electromagnetics-solution-manual-edition-hardcover-by-balanis-constantine-a-published-by-wiley.pdf
- **advanced engineering electromagnetics solutions manual**
  URL: http://denniswildberger.com/advanced-engineering-electromagnetics-solutions-manual.pdf
- **advanced engineering electromagnetics wiley 1989 grading**
  URL: http://denniswildberger.com/advanced-engineering-electromagnetics-wiley-1989-grading.pdf
- **advanced engineering fluid mechanics by biswas**
  URL: http://denniswildberger.com/advanced-engineering-fluid-mechanics-by-biswas.pdf

Tags: #nanda #nursing #diagnosis #chronic #kidney #disease